

08/913218

ABSTRACT

Bz *Compound having a*
A titanium catalyst for reaction between a
carbon-carbon unsaturated bond and a compound having an
5 electrophilic functional group or an electrophilic reagent,
said titanium catalyst being composed of a titanium compound
represented by the formula (1) below



10 (where X^1 , X^2 , X^3 , and X^4 denote independently a halogen atom,
 C_{1-20} alkoxyl group, aralkyloxy group, aryloxy group, or $-\text{NRxRy}$
group (where Rx and Ry denote independently a C_{1-20} alkyl group
or aralkyl group), and any two of X^1 , X^2 , X^3 , and X^4 may form a
ring.) and a Grignard reagent represented by the formula (2)
below in a molar amount 1.5-2.5 times as much as the titanium
15 compound.



(where R^1 denotes a C_{2-10} alkyl group having a hydrogen atom at
the β position and X^5 denotes a halogen atom.)

a
20 The titanium catalyst of the present invention
activates the carbon-carbon unsaturated bond, which has a
comparatively low reactivity, thereby catalyzing the reaction
with an electrophilic functional group. It is inexpensive
and industrially advantageous. When applied to reaction
compound having a
between a carbon-carbon unsaturated bond and an electrophilic
25 functional group, it yields industrially a variety of adducts
of a compound having a carbon-carbon unsaturated bond and a
compound having an electrophilic functional group, and it
also yields a variety of intramolecular adducts of a compound
having a carbon-carbon unsaturated bond and an electrophilic
30 functional group in the same molecule.